Molecular Brain Research, 8 (1990) 359–360
Elsevier

Author Index

Abdahllah, E.A.M., Forray, C. and El-Fakahany, E.E., Relationship between the partial inhibition of muscarinic receptor-mediated phosphoinositide hydrolysis by phorbol esters and tetrodotoxin in rat cerebral cortex, 1
Abood, M.E., Eberwine, J.H., Erdelyi, E. and Evans, C.J., Regulation of both preproenkephalin mRNA and its derived opioids by haloperidol — a method for measurement of peptides and mRNA in the same tissue extract, 243
Abraham, W.C., see Jefferey, K.H., 267
Andrews, G., see Voogt, J.L., 55
Angeretti, N., see Bendotti, C., 343
Anglister, L., see Eichler, J., 213
Arbogast, L.A., see Voogt, J.L., 55

Baird, J.G., see Lambert, D.G., 263
Baker, H., see Stone, D.N., 291
Barnard, E.A., see Fraser, S.P., 331
Battey, J., see Hara, Y., 319
Beaulieu, C., see Jia, W., 311
Beckstead, R.M., see Somers, D.L., 143
Bendotti, C., Servadio, A., Forloni, G., Angeretti, N. and Samanin, R., Increased tryptophan hydroxylase mRNA in raphe serotonergic neurons spared by 5,7-dihydroxytryptamine, 343
Benowitz, L.I., Rodriguez, W.R. and Neve, R.L., The pattern of GAP-43 immunostaining changes in the rat hippocampal formation during reactive synaptogenesis, 17
Berry, R.W., see Redman, R.S., 99
Beyreuther, K., see Malherbe, P., 199
Blake, M.J., Nowak Jr., T.S. and Holbrook, N.J., In vivo hyperthermia induces expression of HSP70 mRNA in brain regions controlling the neuroendocrine response to stress, 89
Blasquez, C., see Bunel, D.T., 167
Bohler Jr., H.C.L., Zoeller, R.T., King, J.C., Rubin, B.S., Weber, R. and Merriam, G.R., Corticotropin releasing hormone mRNA is elevated on the afternoon of proestrus in the parvocellular paraventricular nuclei of the female rat, 259
Bond, R.W., see Wyborski, R.J., 193
Bunel, D.T., Delbende, C., Blasquez, C., Jegou, S. and Vaudry, H., Effects of ions and ionic channel activators or blockers on release of α-MSH from perifused rat hypothalamic slices, 167
Burgoyne, R.D., see Cambray-Deakin, M.A., 77
Burke, W., see Coker III, G.T., 93

Calas, A., see Trembleau, A., 37
Cambray-Deakin, M.A. and Burgoyne, R.D., The non-tyrosinated Mα4α-tubulin gene product is post-translationally tyrosinated in adult rat cerebellum, 77
Chastrette, N., see McCabe, J.T., 325
Cho, W.K., see Lee, B.J., 185
Christakos, S., see Iacopino, A.M., 283
Chuang, D.-M., see Duchemin, A.-M., 235
Coker III, G.T., Studelska, D., Harmon, S., Burke, W. and O'Malley, K.L., Analysis of tyrosine hydroxylase and insulin transcripts in human neuroendocrine tissues, 93
Crawford, M.L.A. and Young, J.M., Ca^{2+}-dependence provides evidence for differing mechanisms of GABA-induced inositol phosphate formation and GABA potentiation of inositol phosphate formation induced by noradrenaline in rat cerebral cortex, 181
Cray, C., Keane, R.W., Malek, T.R. and Levy, R.B., Regulation and selective expression of Ly-6A/E, a lymphocyte activation molecule, in the central nervous system, 9
Cynader, M.S., see Jia, W., 311

Darlison, M.G., see Fraser, S.P., 331
Day, M., see Mack, K., 177
Delbende, C., see Bunel, D.T., 167
Djamgoz, M.B.A., see Fraser, S.P., 331
Dragunow, M., see Jefferey, K.J., 267
Draguhn, A., see Malherbe, P., 199
Duchemin, A.-M., Quach, T.T., Schrier, B.K., Chuang, D.-M. and Wyatt, R.J., Expression of neurotrophic activity in *Xenopus* oocytes injected with mRNA from wounded rat cerebral cortex, 235

Eberwine, J.H., see Abood, M.E., 243
Eichler, J., Silman, I., Gentry, M.K. and Anglister, L., Immunocytochemical localization of phosphatidylinositol-anchored acetylcholinesterase in excitable membranes of *Torpedo ocellata*, 213
El-Fakahany, E.E., see Abdallah, E.A.M., 1
Erdelyi, E., see Abood, M.E., 243
Evans, C.J., see Abood, M.E., 243

Fader, D. and Lewis, E.J., Interaction of cyclic AMP and cell-cell contact in the control of tyrosine hydroxylase RNA, 25
Fitzgerald, L.W., see Ratty, A.K., 355
Forloni, G., see Bendotti, C., 343
Forray, C., see Abdallah, E.A.M., 1
Fraser, S.P., Djamgoz, M.B.A., Usherwood, P.N.R., O'Brien, J., Darlison, M.G. and Barnard, E.A., Amino acid receptors from insect muscle: electrophysiological characterization in *Xenopus* oocytes following expression by injection of mRNA, 331
Fèvre-Montange, M., see Trembleau, A., 37

Gainer, H., see Hara, Y., 319
Ganten, D., see McCabe, J.T., 325
Gentry, M.K., see Eichler, J., 213
Giros, B., see Llorens-Cortes, C., 227
Glick, S.D., see Ratty, A.K., 355
Gottlieb, D.E., see Mack, K., 177
Gottlieb, D.I., see Wyborski, R.J., 193
Grasso, A., see Umbach, J.A., 31
Gross, K.W., see Ratty, A.K., 355
Gubits, R.M., Wollack, J.B., Yu, H. and Liu, W.-K., Activation of adenosine receptors induces c-*fos*, but not c-*jun*, expression in neuron-glia hybrids and fibroblasts, 275
Gundersen, C.B., see Umbach, J.A., 31

Hall, A.K., Hempstead, J. and Morgan, J.I., Thymosin 10 levels in developing human brain and its regulation by retinoic acid in the HTB-10 neuroblastoma, 129
Hara, Y., Battey, J. and Gainer, H., Structure of mouse vasopressin and oxytocin genes, 319
Harmon, S., see Coker III, G.T., 93
Harry, J., see Stahl, N., 209
Hart, R.P., see Walker, P.D., 113
Hempstead, J., see Hall, A.K., 129
Hicks, K.J., see Hisanaga, K., 69
Hisanaga, K., Sagar, S.M., Hicks, K.J., Swanson, R.A. and Sharp, F.R., c-*fos* proto-oncogene expression in astrocytes associated with differentiation or proliferation but not depolarization, 69
Holbrook, N.J., see Blake, M.J., 89
Hong, J.S., see Li, S.J., 219
Howells, R.D., see Lauber, A.H., 47
Huang, F.L., see Jia, W., 311
Hudson, P.M., see Li, S.J., 219
Hänze, J., see McCabe, J.T., 325

Iacopino, A.M., Rhoten, W.B. and Christakos, S., Calcium binding protein (calbindin-D_{28k}) gene expression in the developing and aging mouse cerebellum, 283

Jacobowitz, D.M., see Sutin, E.L. 63
Jefferey, K.J., Abraham, W.C., Dragunow, M. and Mason, S.E., Induction of Fos-like immunoreactivity and the maintenance of long-term potentiation in the dentate gyrus of unanesthetized rats, 267
Jegou, S., see Bunel, D.T., 167
Jia, W., Beaulieu, C., Huang, F.L. and Cynader, M.S., Cellular and subcellular localization of protein kinase C in cat visual cortex, 311
Jiang, H.K., see Li, S.J., 219
Joh, T.H., see Kang, U.J., 83
Joh, T.H., see Stone, D.N., 291
Jonakait, G.M., see Walker, P.D., 113

Kang, U.J. and Joh, T.H., Deduced amino acid sequence of bovine aromatic L-amino acid decarboxylase: homology to other decarboxylase, 83
Keane, R.W., see Cray, C., 9
Kikuchi, A., see Yamamoto, J., 105
Kilbourne, E.J. and Sabban, E.L., Differ-

ential effect of membrane depolarization on levels of tyrosine hydroxylase and dopamine β-hydroxylase mRNAs in PC12 pheochromocytoma cells, 121
Kim, K., see Lee, B.J., 185
King, J.C., see Bohler Jr., H.C.L., 259

Lambert, D.G., Whitham, E.M., Baird, J.G. and Nahorski, S.R., Different mechanisms of Ca^{2+} entry induced by depolarization and muscarinic receptor stimulation in SH-SY5Y human neuroblastoma cells, 263
Lang, R.E., see McCabe, J.T., 325
Lauber, A.H., Romano, G.J., Mobbs, C.V., Howells, R.D. and Pfaff, D.W., Estradiol induction of proenkephalin messenger RNA in hypothalamus: dose–response and relation to reproductive behavior in the female rat, 47
Lee, B.J., Kim, K. and Cho, W.K., Activation of intracellular pathways with forskolin and phorbol ester increases LHRH mRNA level in the rat hypothalamus superfused in vitro, 185
Lehmann, E., see McCabe, J.T., 325
Levy, R.B., see Cray, C., 9
Lewis, E.J., see Fader, D., 25
Li, S.J., Jiang, H.K., Stachowiak, M.S., Hudson, P.M., Owyang, V., Nanry, K., Tilson, H.A. and Hong, J.S., Influence of nigrostriatal dopaminergic tone on the biosynthesis of dynorphin and enkephalin in rat striatum, 219
Liu, W.-K., see Gubits, R.M., 275
Llorens-Cortes, C., Van Amsterdam, J.G.C., Giros, B., Quach, T.T. and Schwartz, J.C., Enkephalin biosynthesis and release in mouse striatum are inhibited by GABA receptor stimulation: compared changes in preproenkephalin mRNA and Tyr-Gly-Gly levels, 227

Mack, K., Day, M., Milbrandt, J. and Gottlieb, D.E., Localization of the NGFI-A protein in the rat brain, 177
Malek, T.R., see Cray, C., 9
Malherbe, P., Draguhn, A., Multhaup, G., Beyreuther, K. and Möhler, H., $GABA_A$-receptor expressed from rat brain α- and β-subunit cDNAs displays potentiation by benzodiazepine receptor ligands, 199
Mason, S.E., see Jefferey, K.J., 267
Maul, G.G., see Reddy, U.R., 137
McCabe, J.T., Lehmann, E., Chastrette, N., Hänze, J., Lang, R.E., Ganten, D. and Pfaff, D.W., Detection of vasopressin mRNA in the neurointermediate lobe of the rat pituitary, 325
Merriam, G.R., see Bohler Jr., H.C.L., 259
Micevych, P.E., see Popper, P., 159
Milbrandt, J., see Mack, K., 177
Mobbs, C.V., see Lauber, A.H., 47
Morgan, J.I., see hall, A.K., 129
Mullins, J.J., see Ratty, A.K., 355
Multhaup, G., see Malherbe, P., 199
Möhler, H., see Malherbe, P., 199

Nahorski, S.R., see Lambert, D.G., 263
Nanry, K., see Li, S.J., 219
Neve, R.L., see Benowitz, L.I., 17
Nowak Jr., T.S., see Blake, M.J., 89

O'Brien, J., see Fraser, S.P., 331
O'Malley, K.L., see Coker III, G.T., 93
Ohga, N., see Yamamoto, J., 105
Okamoto, K., see Sekiguchi, M., 301
Owyang, V., see Li, S.J., 219

Pfaff, D.W., see Lauber, A.H., 47
Pfaff, D.W., see McCabe, J.T., 325
Phillips, L.L. and Steward, O., Increases in mRNA for cytoskeletal proteins in the denervated neuropil of the dentate gyrus: an in situ hybridization study using riboprobes for β-actin and β-tubulin, 249
Popko, B., see Stahl, N., 209
Popper, P. and Micevych, P.E., Steroid regulation of calcitonin gene-related peptide mRNA expression in motoneurons of the spinal nucleus of the bulbocavernosus, 159

Quach, T.T., see Duchemin, A.-M., 235
Quach, T.T., see Llorens-Cortes, C., 227
Quadri, S.K., see Voogt, J.L., 55

Ratty, A.K., Fitzgerald, L.W., Titeler, M., Glick, S.D., Mullins, J.J. and Gross, K.W., Circling behavior exhibited by a transgenic insertional mutant, 355
Reddy, U.R., Venkatakrishnan, G., Maul, G.G., Roy, A.K. and Ross, A.H., Transient expression of full-length and truncated forms of the human nerve growth factor receptor, 137
Redman, R.S. and Berry, R.W. α-Bag cell peptide inhibits bag cell adenylate cyclase via a GTP-dependent mechanism, 99
Rhoten, W.B., see Iacopino, A.M., 283
Rodriguez, W.R., see Benowitz, L.I., 17
Romano, G.J., see Lauber, A.H., 47
Ross, A.H., see Reddy, U.R., 137
Roy, A.K., see Reddy, U.R., 137
Rubin, B.S., see Bohler Jr., H.C.L., 259

Sabban, E.L., see Kilbourne, E.J., 121
Sagar, S.M., see Hisanaga, K., 69
Sakai, Y., see Sekiguchi, M., 301
Sakuta, H., see Sekiguchi, M., 301
Samanin, R., see Bendotti, C., 343
Schotland, S., see Walker, P.D., 113
Schrier, B.K., see Duchemin, A.-M., 235
Schwartz, J.C., see Llorens-Cortes, C., 227
Sekiguchi, M., Sakuta, H., Okamoto, K. and Sakai, Y., $GABA_B$ receptors expressed in *Xenopus* oocytes by guinea pig cerebral mRNA are functionally coupled with Ca^{2+}-dependent Cl^- channels and with K^+ channels, through GTP-binding proteins, 301
Senba, E., Simmons, D.M., Wada, E., Wada, K. and Swanson, L.W., RNA levels of neuronal nicotinic acetylcholine receptor subunits are differentially regulated in axotomized facial motoneurons: an in situ hybridization study, 349
Servadio, A., see Bendotti, C., 343
Sharp, F.R., see Hisanaga, K., 69
Silman, I., see Eichler, J., 213
Simmons, D.M., see Senba, E., 349
Somers, D.L. and Beckstead, R.M., Striatal preprotachykinin and preproenkephalin mRNA levels and the levels of nigral substance P and pallidal Met^5-enkephalin depend on corticostriatal axons that use the excitatory amine acid neurotransmitters aspartate and glutamate: quantitative radioimmunocytochemical and in situ hybridization evidence, 143
Stachowiak, M.S., see Li, S.J., 219
Stahl, N., Harry, J. and Popko, B., Quantitative analysis of myelin protein gene expression during development in the rat sciatic nerve, 209
Steward, O., see Phillips, L.L., 249
Stone, D.N., Wessel, T., Joh, T.H. and Baker, H., Decrease in tyrosine hydroxylase, but not aromatic L-amino acid decarboxylase, messenger RNA in rat olfactory bulb following neonatal, unilateral odor deprivation, 291
Studelska, D., see Coker III, G.T., 93
Sutin, E.L. and Jacobowitz, D.M., Detection of CCK mRNA in the motor nucleus of the rat trigeminal nerve with in situ hybridization histochemistry, 63
Swanson, L.W., see Senba, E, 349
Swanson, R.A., see Hisanaga, K., 69

Takai, Y., see Yamamoto, J., 105
Tilson, H.A., see Li, S.J., 219
Titeler, M., see Ratty, A.K., 355
Trembleau, A., Calas, A. and Fèvre-Montange, M., Ultrastructural localization of oxytocin mRNA in the rat hypothalamus by in situ hybridization using a synthetic oligonucleotide, 37

Ueda, T., see Yamamoto, J., 105
Umbach, J.A., Grasso, A. and Gundersen, C.B., α-Latrotoxin triggers an increase of ionized calcium in *Xenopus* oocytes injected with rat brain mRNA, 31
Usherwood, P.N.R., see Fraser, S.P., 331

Van Amsterdam, J.G.C., see Llorens-Cortes, C., 227
Vaudry, H., see Bunel, D.T., 167
Venkatakrishnan, G., see Reddy, U.R., 137
Voogt, J.L., Arbogast, L.A., Quadri, S.K. and Andrews, G., Tyrosine hydroxylase messenger RNA in the hypothalamus, substantia nigra and adrenal medulla of old female rats, 55

Wada, E., see Senba, E., 349
Wada, K., see Senba, E., 349
Walker, P.D., Schotland, S., Hart, R.P. and Jonakait, G.M., Tryptophan hydroxylase inhibition increases preprotachykinin mRNA in developing and adult medullary raphe nuclei, 113
Weber, R., see Bohler Jr., H.C.L., 259
Wessel, T., see Stone, D.N., 291
Whitham, E.M., see Lambert, D.G., 263
Wollack, J.B., see Gubits, R.M., 275
Wyatt, R.J., see Duchemin, A.-M., 235
Wyborski, R.J., Bond, R.W. and Gottlieb, D.I., Characterization of a cDNA coding for rat glutamic acid decarboxylase, 193

Yamamoto, J., Kikuchi, A., Ueda, T., Ohga, N. and Takai, Y., A GTPase-activating protein for *rho*B p20, a *ras* p21-like GTP-binding protein — partial purification, characterization and subcellular distribution in rat brain, 105
Young, J.M., see Crawford, M.L.A., 181
Yu, H., see Gubits, R.M., 275

Zoeller, R.T., see Bohler Jr., H.C.L., 259